

drug	stop
ACTONEL[RISEDRONATE]	STOP 3 DAYS BEFORE SURGERY, RESUME 3 DAYS LATER--WHEN ANTIBIOTICS FINISHED
<i>ASPIRIN</i>	<i>STOP 5-7 DAYS BEFORE SURGERY</i>
BRILINTA [TICAGRELOR]	STOP 5 DAYS BEFORE SURGERY
<i>CELEBREX [CELECOXIB]</i>	<i>STOP 2 DAYS BEFORE SURGERY</i>
CLINORIL [SULINDAC]	STOP 3 DAYS BEFORE SURGERY
<i>COUMADIN [WARFARIN]</i>	<i>STOP 5 DAYS BEFORE SURGERY</i>
DISEASE MODIFYING ANTI RHEUMATIC DRUGS	STOP 1-2 WEEKS BEFORE SURGERY, RESUME 1-2 WEEKS AFTER SURGERY
<i>DOLOBID [DIFLUNISAL]</i>	<i>STOP 3 DAYS BEFORE SURGERY</i>
EFFIENT [PRASUGREL]	STOP 7 DAYS BEFORE SURGERY
<i>ELIQUIS [APIXABAN]</i>	<i>STOP 2 DAYS BEFORE SURGERY</i>
ENBREL [ETANERCEPT]	STOP 2 WEEKS BEFORE SURGERY, RESUME 2 WEEKS AFTER
<i>ESTROGEN / BIRTH CONTROL PILLS</i>	<i>?? STOP ONE MONTH BEFORE SURGERY?? WEIGH RISK vs. BENEFIT</i>
FELDENE [PIROXICAM]	STOP 10 DAYS BEFORE SURGERY
<i>FOSAMAX</i>	<i>STOP AT TIME OF SURGERY</i>
GLUCOPHAGE	STOP 24 HOURS BEFORE SURGERY
<i>HERBALS AND SUPPLEMENTS</i>	<i>STOP ONE WEEK BEFORE SURGERY</i>
HUMIRA [ADLAIMUBAB]	DO SURGERY DURING THE WEEK THAT PATIENT DOES NOT GET AN INJECTION
<i>IMURAN [AZATHIOPRINE]</i>	<i>STOP 3 DAYS BEFORE SURGERY--RESUME 3 DAYS AFTER</i>
INDOCIN [INDOMETHACIN]	STOP ONE DAY BEFORE SURGERY
<i>MAO INHIBITORS --SELIGILINE , RASIGILINE</i>	<i>TAPER OFF 2-3 WEEKS BEFORE PROCEDURE</i>
METHOTREXATE	<i>OK TO CONTINUE UNLESS CO-MORBIDITIES. OTHERWISE 1 WEEK BEFORE>1 AFTER</i>
<i>MOBIC [MELOXICAM]</i>	<i>STOP 10 DAYS BEFORE SURGERY</i>
MOTRIN [IBUPROFEN]	STOP ONE DAY BEFORE SURGERY
<i>NAPROSYN [NAPROXEN]</i>	<i>STOP 3 DAYS BEFORE SURGERY</i>
ORUDIS [KETOPROFEN]	STOP ONE DAY BEFORE SURGERY
<i>PERSANTINE [DIPYRIMADOLE]</i>	<i>STOP 2 DAYS BEFORE SUGERY</i>
PLAQUENIL [HYDROXYCHLOROQUINE]	STOP 3 DAYS BEFORE SURGERY--RESUME 3 DAYS AFTER
<i>PLAVIX [CLOPIDOGREL]</i>	<i>STOP 5 DAYS BEFORE SURGERY</i>
PLETAL [CILOSTAZOL]	STOP 3 DAYS BEFORE SURGERY

<i>PRADAXA [DABIGATRAN]</i>	<i>STOP 2 DAYS BEFORE IF CrCl > 50--STOP 5 DAYS BEFORE IF Cr Cl < 50</i>
RELAFEN [NABUMETONE]	STOP 10 DAYS BEFORE SURGERY
<i>REMICADE [INFLIXIMAB]</i>	<i>STOP 4 WEEKS BEFORE SURGERY, RESUME 4 WEEKS AFTER</i>
SINEMET	DO NOT STOP
<i>TAMOXIFEN / RALOXIFENE</i>	<i>STOP ONE WEEK BEFORE SURERY, IF PROCEDURE IS HIGH RISK FOR DVT</i>
TICLID [TICLOPIDINE]	STOP 5 DAYS BEFORE SURGERY
<i>TNF [TUMOR NECROSIS FACTOR] DRUGS</i>	<i>STOP 1-2 WEEKS BEFORE SURGERY, RESUME 1-2 WEEKS AFTER SURGERY</i>
VOLTAREN [DICLOFENAC]	STOP ONE DAY BEFORE SURGERY
<i>XARELTO [RIVAROXABAN]</i>	<i>STOP 2 DAYS BEFORE SURGERY</i>